

EXHIBIT A

Lasers in Cutaneous and Aesthetic Surgery

EDITORS

Kenneth A. Arndt, MD

Professor of Dermatology
Harvard Medical School
Boston, Massachusetts
Dermatologist-in-Chief
Codirector, Joint Center for Cutaneous Laser Surgery
Beth Israel Deaconess Medical Center
Boston, Massachusetts

Jeffrey S. Dover, MD

Associate Professor of Clinical Dermatology
Harvard Medical School
Boston, Massachusetts
Codirector, Joint Center for Cutaneous Laser Surgery
Department of Dermatology
Beth Israel Deaconess Medical Center
Boston, Massachusetts

Suzanne M. Olbricht, MD

Assistant Professor of Clinical Dermatology
Harvard Medical School
Boston, Massachusetts
Director of Dermatology Surgery
Department of Dermatology
Beth Israel Deaconess Medical Center
Boston, Massachusetts



Lippincott - Raven

P U B L I S H E R S

Philadelphia • New York

Manufacturing Manager: Dennis Teston
Associate Managing Editor: Kathleen Bubbeo
Cover Designer: Karen Quigley
Production Service: Textbook Writers Associates, Inc.
Indexer: Michael Loo
Compositor: Compset Inc.
Printer: Kingsport Press

©1997, by Lippincott-Raven Publishers. All rights reserved. This book is protected by copyright. No part of it may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means—electronic, mechanical, photocopy, recording, or otherwise—without the prior written consent of the publisher, except for brief quotations embodied in critical articles and reviews. For information write Lippincott-Raven Publishers, 227 East Washington Square, Philadelphia, PA 19106-3780.

Materials appearing in this book prepared by individuals as part of their official duties as U.S. Government employees are not covered by the above-mentioned copyright.

Printed in the United States of America

987654321

Library of Congress Cataloging-in-Publication Data
Lasers in cutaneous and aesthetic surgery/editors, Kenneth A. Arndt,
Jeffrey S. Dover, Suzanne M. Olbricht.

p. cm.

Includes bibliographical references and index.

ISBN 0-316-05177-2

I. Skin—Laser surgery. 2. Surgery, Plastic. 3. Lasers in surgery. I. Arndt, Kenneth A., 1936—. II. Dover, Jeffrey S. III. Olbricht, Suzanne M.

[DNLM: 1. Laser Surgery—methods. 2. Skin—surgery. 3. Surgery, Plastic—methods. WO 511 L3436 1997]

RL120.L37L357 1997

617.4'77059—dc21

DNLM/DLC

for Library of Congress

97-3099

CIP

Care has been taken to confirm the accuracy of the information presented and to describe generally accepted practices. However, the authors, editors, and publisher are not responsible for errors or omissions or for any consequences from application of the information in this book and make no warranty, express or implied, with respect to the contents of the publication.

The authors, editors, and publisher have exerted every effort to ensure that drug selection and dosage set forth in this text are in accordance with current recommendations and practice at the time of publication. However, in view of ongoing research, changes in government regulations, and the constant flow of information relating to drug therapy and drug reactions, the reader is urged to check the package insert for each drug for any change in indications and dosage and for added warnings and precautions. This is particularly important when the recommended agent is a new or infrequently employed drug.

Some drugs and medical devices presented in this publication have Food and Drug Administration (FDA) clearance for limited use in restricted research settings. It is the responsibility of the health care provider to ascertain the FDA status of each drug or device planned for use in their clinical practice.

for all photobiologic effects and laser-tissue interactions to occur. The

This chapter was adapted from Anderson RR. Laser tissue interactions. In: Goldman MP, Fitzpatrick RE, eds. Cutaneous laser surgery. 1994. St. Louis: Mosby-Year Book, 1994.

injury occurs. The relatively low power continuous-wave lasers such as the CO₂ and argon-ion laser and the quasi-continuous-wave (rapid-pulsed) lasers such as the copper vapor and KTP lasers usually cause a well-controlled, superficial, partial-thickness burn. In contrast, pulsed yellow-dye lasers designed for selective photothermolysis of microvascular lesions selectively coagulate microvessels.

Selective photothermolysis is a term coined by Dr. Parrish and myself¹⁰ over 14 years ago to describe the sequence of light-heat-ruin that occurs at microscopic sites, such as blood vessels, pigmented cells, and tattoo ink particles, which selectively absorb light pulses. Selective photothermolysis is qualitatively different from gross thermal injury and is like a "magic bullet" in its ability to target certain structures. Short pulses are necessary to deposit energy in the targets before they can cool off, thus achieving extreme, localized heating of the targets. Thermal coagulation or thermally mediated mechanical damage, or both, are involved, depending on the rate of energy deposition in the targets. This technique led to the development of a new generation of dermatologic lasers that cause much less scarring, and these are discussed in detail here.

Mechanical injury, sometimes misnamed "photoacoustic injury," occurs as the result of the rapid heating produced by high-energy, short-pulse lasers. The rate of local heating can be so severe that structures are torn apart by *shock waves* (a highly destructive supersonic pressure wave), *cavitation* (the sudden expansion and collapse of a steam bubble), or rapid *thermal expansion*. Terms such as *fracture*, *rupture*, or *explosion* are appropriately used in reference to these processes, even though they occur on a microscopic scale. Mechanical damage plays a central, positive role in the tattoo removal achieved by selective photothermolysis using high-energy, submicrosecond lasers. However, mechanical damage can also be an unwanted side effect, such as in the form of the bleeding caused by pulsed yellow-dye lasers and the ejection of tissue fragments caused by high-energy, Q-switched laser pulses.

THERMAL INJURY TO CELLS

Most human cells can withstand prolonged exposure to a temperature of 40°C. At 45°C, cultured human fibroblasts have been observed to be destroyed after about 20 minutes of exposure. However, the same cells can withstand a temperature of more than 100°C if exposed to it for only 10⁻³ seconds.¹¹ Thus, it is not the temperature per se but a combination of the local temperature and the heating time that determines whether cells in the skin are injured. This is presumably because thermal denaturation is a rate-dependent process, as described earlier. This behavior has an important bearing on the cell injury that occurs in the setting of selective photothermolysis, during which extreme temperatures are present in target sites, such as microvessels, for a very short duration. Essentially nothing is known

Light deposits energy only at sites of absorption. Heat is created at wavelengths that penetrate into skin and are preferentially absorbed by chromophoric structures, such as blood vessels or melanin-containing cells. As soon as the heat is created, however, it begins to dissipate as the result of conduction and radiative transfer. Thus, the competition between active heating and passive cooling determines how hot the targets become. The most selective target heating is achieved when the energy is deposited at faster than the rate of cooling of the target structures.

Three basic elements are necessary to achieve selective photothermolysis: (1) a wavelength that reaches and is preferentially absorbed by the desired target structures, (2) an exposure duration less than or equal to the time necessary for cooling of the target structures, and (3) sufficient fluence to produce a damaging temperature in the target structures. When these criteria are met, exquisitely selective injury occurs in thousands of microscopic targets, without the need to aim the laser at each one. The effect is equivalent to the legendary magic bullets that seek out only the desired target. Selective photothermolysis makes use of a variety of thermally mediated damage mechanisms, including thermal denaturation, mechanical damage resulting from rapid thermal expansion or phase changes (cavitation), and pyrolysis (changes in the primary chemical structure).

A useful construct is the *thermal relaxation time*, mentioned earlier. When the laser exposure duration is less than the thermal relaxation time, maximal thermal confinement occurs because the target cannot then get rid of its heat during the laser exposure. Many processes are involved in cooling, including convection, radiation, and conduction. Of these, thermal conduction is the one that dominates the cooling of microscopic structures in skin. However, microscale radiational cooling in tissue has never been thoroughly examined and in theory may be of importance for very small targets at high temperatures, such as tattoo ink particles or melanin granules.

It is common experience that smaller objects cool more quickly than larger objects. For example, a cup of tea cools faster than a hot bath, even though both involve hot water in a porcelain container. More precisely, the thermal relaxation time for heat conduction is proportional to the square of size (see Eq. 2). That is, for any given material and shape, an object half the size will cool in one fourth the time and an object one tenth the size will cool in one hundredth the time. This behavior is important in optimizing the pulse duration or the exposure duration for the selective photothermolysis of blood vessels. Blood vessels vary in size, from capillaries, which have thermal relaxation times of tens of microseconds, to venules and arterioles, which have thermal relaxation times of hundreds of microseconds, to the large venules of adult PWSs, which have thermal relaxation times of tens to hundreds of milliseconds. This means that there are vessels in a typical adult PWS with thermal relaxation times ranging over three orders of magnitude, and thus it is ludicrous to attempt to define "the" thermal relaxation time for vessels.

Lasers and
lasers in
with an ab-
about 200
of the CO₂
hind about
tered in less
oped mainly
ed for endo-
hemostasis.
quiring ex-

strongly ab-
high precision,
of about 10,000
for tissue re-
delivered in a few
per pulse, leaving
nm. Short-pulsed
only one or two cell
y. This is excellent
or choice when he-
achieved (and ther-
nger pulses, which
o or other lasers op-
d should therefore
not yet been devel-

t, have been shown
photochemical ab-
ent is about 12,000
or the erbium laser
ent to break chem-
not only through
hrough the photo-
late, excimer laser
ses, other than the
however, because of
te the cornea, this
l refraction.¹⁹ The
ation of skin, pro-
ill into the epider-

pe of the uses of
elopment.¹⁰ It is by

A service of the U.S. National Library of Medicine and the National Institutes of Health

My NCBI
[Sign In] [Register]

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC
Journals Books

Search PubMed for

Go Clear

Advanced Search (beta)

Limits Preview/Index History Clipboard Details

Display AbstractPlus

Show 20

Sort By

Send to

All: 1

Review: 0

1: J Invest Dermatol. 1983 Jun;80 Suppl:75s-80s.

Links

Selective thermal effects with pulsed irradiation from lasers: from organ to organelle.

Parrish JA, Anderson RR, Harrist T, Paul B, Murphy GF.

Specific damage by selectively absorbed, pulsed lasers can be predicted based on physical models. Thermally mediated alterations can be confined to pigmented targets from the level of subcellular organelles (e.g., melanosomes) to large multicellular tissue structures (e.g., blood vessels) by the appropriate manipulation of wavelength and pulse duration. Highly selective damage to human cutaneous microvessels in vivo is shown to occur after 0.3-microseconds 577-nm dye laser pulses; the epidermis and dermal structures other than vessels are spared. Observations in an animal model suggest that hemorrhage or, at lower doses, selective intravascular coagulation and permanent microvascular hemostasis occur. Highly selective damage to melanized cells and to single melanosomes in situ was shown to occur after single 20-ns 351-nm pulses from a XeF excimer laser. Basal-cell- and melanocyte-specific necrosis is followed by gross hypopigmentation. In this case there is no evidence of vascular damage. The most likely modes of selective alterations include localized thermal denaturation, vaporization, and shock-wave generation. Means of predicting and controlling histologically selective radiant heating effects in skin are suggested.

PMID: 6854059 [PubMed - indexed for MEDLINE]

Related Articles

Effect of dye laser pulse duration on selective cutaneous microvascular injury. [J Invest Dermatol. 1986]

Microvasculature can be selectively damaged using dye lasers: a basic theory and experimental evidence in human skin. [Lasers Surg Med. 1981]

Mechanisms of selective vascular changes caused by dye lasers. [Lasers Surg Med. 1983]

Organelle-specific injury to melanin-containing cells in human skin by pulsed laser irradiation. [Lab Invest. 1983]

Ultrastructure: effects of melanin pigment on target specificity using a pulsed dye laser (577 nm). [J Invest Dermatol. 1987]

» See all Related Articles ..

Display AbstractPlus

Show 20

Sort By

Send to

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Privacy Statement | Freedom of Information Act | Disclaimer

A service of the U.S. National Library of Medicine
and the National Institutes of Health

My NCBI
[Sign In] [Register]

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC
Journals Books

Search PubMed for

Go

Clear

Advanced Search (beta)

Limits Preview/Index History Clipboard Details

Note: Performing your original search, *selective photothermolysis*, in PubMed will retrieve 137 citations.

Display AbstractPlus

Show 20

Sort By

Send to

All: 1 Review: 0

1: Science. 1983 Apr 29;220(4596):524-7.

Science

Links

Selective photothermolysis: precise microsurgery by selective absorption of pulsed radiation.

Anderson RR, Parrish JA.

Suitably brief pulses of selectively absorbed optical radiation can cause selective damage to pigmented structures, cells, and organelles in vivo. Precise aiming is unnecessary in this unique form of radiation injury because inherent optical and thermal properties provide target selectivity. A simple, predictive model is presented. Selective damage to cutaneous microvessels and to melanosomes within melanocytes is shown after 577-nanometer (3×10^{-7} second) and 351-nanometer (2×10^{-8} second) pulses, respectively. Hemodynamic, histological, and ultrastructural responses are discussed.

PMID: 6836297 [PubMed - indexed for MEDLINE]

Related Articles

Selective thermal effects with pulsed irradiation from lasers: from organ to organelle. [J Invest Dermatol. 1983]

Non-invasive laser microsurgery selectively damages populations of labeled mouse neurons: dependence on incident laser dose and [J Neurosci. 1988]

Melanin granule models for the processes of laser-induced thermal damage in pigmented retinal tissues. I. Modeling of laser-induced healing of melanosomes and selective thermal processes in retinal [Regul Mech Biol. 2007]

Ex vivo quality-switched ruby laser irradiation of cutaneous melanocytic lesions: persistence of S-100-, HMB-45- and Masson-positive [Dermatol Surg. 1997]

Selective targeting of trabecular meshwork cells: in vitro studies of pulsed and CW laser interaction [Invest Ophthalmol Vis Sci. 1995]

» See all Related Articles..

Display AbstractPlus

Show 20

Sort By

Send to

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Privacy Statement | Freedom of Information Act | Disclaimer